



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

UCRL-CONF-202979

# **The Kinematics of the Lag-Luminosity Relationship**

*Jay D. Salmonson*

**December 1, 2002**

Submitted to The 20<sup>th</sup> Texas Symposium on Relativistic Astrophysics on December 10-15, 2002 in Austin, Texas.

## **Disclaimer**

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

# The Kinematics of the Lag-Luminosity Relationship

Jay D. Salmonson

*Lawrence Livermore National Laboratory, Livermore, California 94550*

**Abstract.** Herein I review the argument that kinematics, i.e. relativistic motions of the emitting source in gamma-ray bursts (GRBs), are the cause of the lag-luminosity relationship observed in bursts with known redshifts.

## INTRODUCTION


Norris *et al.* [1] have discovered a relationship between the peak luminosity of gamma-ray bursts (GRB) and the pulse time lag between BATSE energy channels. In Salmonson 2000 [2] this correlation was shown to be improved when we neglect our poor knowledge of received photon energy. Thus instead we use the correlation between photon number luminosity and pulse time lag. It was found that the inferred isotropic peak number luminosity  $N_{pk}$  (photons  $\text{s}^{-1}$ ) is related to the observed spectral lag between energy channels  $\Delta t$  by

$$N_{pk} = 8.6 \times 10^{56} \Delta t^{-0.98} . \quad (1)$$

I argue that kinematics are the origin of this relation [2]. Specifically, bursts with emitting material moving with a higher velocity toward the observer appear more luminous and have shorter observed lags (derived from an intrinsic pulse cooling timescale) between observed energy channels due to relativistic blue shift. Relativistic beaming allows one to only consider emitters moving directly toward the observer. I propose that the wide range of observed (cosmological redshift compensated) spectral lags and inferred luminosities (see Figure 1) can be explained if GRBs derive from a relativistic jet, with opening angle  $\theta_0$ , in which the fastest material moves along the core of the jet and the velocity of the material monotonically decreases with increasing angle from the jet axis. The variety of observed bursts then derives from our perspective of the jet. All of the material is presumed to move relativistically ( $\gamma \gg 1$ ) and so all of our received flux is derived from a very small  $\sim 1/\gamma^2$  solid angle of the jet; much smaller than the jet opening angle ( $1/\gamma \ll \theta_0$ ). It is from this small region that all of our information about a burst is derived.

This hypothesis that all GRBs derive from a single jet morphology viewed at various orientations then predicts that there will be GRB jets viewed at such high inclinations, and thus with such low Lorentz factors that the relativistic beaming angle will be larger than the jet opening angle ( $1/\gamma > \theta_0$ ). The emission will no longer be consistent with isotropy and thus I predict a break in the lag-luminosity relation:  $N_{pk} \propto \Delta t^{-3}$  [3]. Due to the low Lorentz factor, such bursts would be very dim. If GRB980425, which was unusual in several respects [3] and, by its association with supernova 1998bw [4], was sub-luminous, defines this break (see Figure 1), then one can calculate key features of the GRB jet. In particular, Lorentz factors range from  $\gamma_{\max} \sim 1000$  for bright bursts such as GRB990123, to  $\gamma_{\text{break}} \sim 100$  for more middling bursts like GRB980703. The position of the break implies an opening angle of  $\theta_0 \sim 1^\circ$  and thus a total gamma-ray burst energy of  $E_{\text{tot}} \sim 10^{50}$  ergs, which is much less than isotropic energy estimates (up to  $10^{54}$  ergs [5]).

This work was performed under the auspices of the U.S. Department of Energy by University of California Lawrence Livermore National Laboratory under contract W-7405-ENG-48.



**FIGURE 1.** Peak photon number luminosity  $N_{pk}$  versus spectral pulse lag for six bursts with known redshifts plus GRB980425. A break is inferred by fitting a break slope  $\propto \Delta t^{-3}$  to intersect GRB980425. Spectral cross-correlation function lags between BATSE channels 3 and 1 (CCF31) for regions down to 0.5 and 0.1 of peak intensity were obtained from Norris *et al.* (2000). The line of best fit for 0.1 (squares) is  $\propto \Delta t^{-0.98}$ .

## REFERENCES

1. Norris J. P., Marani G. F., Bonnell J. T., *ApJ* **534**, 248 (2000).
2. Salmonson J. D., *ApJL* **544**, L115 (2000).
3. Salmonson J. D., *ApJL* **546**, L29 (2001).
4. Galama T. J. et al., *Nature* **395**, 670 (1998).
5. Galama T. J. et al., *Nature* **398**, 394 (1999).